

Ground state properties and bubble structure of superheavy nuclei

Shailesh K. Singh¹, Mohammad Ikram² and S. K. Patra¹

¹ *Institute of Physics, Sachivalaya Marg, Bhubaneswar-751005, India*

² *Department of Physics, Aligarh Muslim University, Aligarh-202002, India.*

(Dated: July 13, 2012)

Abstract

We calculate the ground state properties of recently synthesized superheavy nuclei starting from $Z=105$ -120. The nonrelativistic and relativistic mean field formalisms are used to evaluate the binding energy, charge radius, quadrupole deformation parameter and the density distribution of nucleons. We analyzed the stability of the nuclei based on the binding energy and neutron to proton ratio. We also studied the bubble structure of the nucleus which reveals about the special features of the superheavy nucleus.

PACS numbers:

I. INTRODUCTION

The study of superheavy nuclei is an interesting topic for the current days research in nuclear physics. Initially, the transuranium elements were made by subjecting Uranium to a high neutron flux allowing to capture neutrons successively. The neutron-rich isotopes undergo β -decay and produce new elements. This process allows to prepare element maximum upto Fermion ($Z = 100$), because of the shorter time interval for the spontaneous fission of the new element than neutron capture time. To make heavier Z nuclei one has to reach the so called *island of stability* which is a matter of discussion from last five decades [1–4]. Earlier, it was predicted [1, 5] that the next proton magic number would be $Z=126$, because of the equality of the proton and neutron magic numbers for known closed shell nuclei. However, several microscopic calculations [6–11] suggest a shift of this number to 114. One of the cause of the shift is Coulomb effects on the spherical single particle levels. The use of shell correction by V. M. Strutinsky [12] to the liquid-drop calculation of binding energy (BE) opens a more satisfactory exploration of double close nuclei beyond ^{208}Pb . Recently, based on the relativistic mean field (RMF) and Skyrme Hartree-Fock (SHF) calculations Patra and collaborator [13, 14] predicts that the *island of stability* in the superheavy region is situated near mass number $A \sim 300$ peaking at $^{292,304}120$.

Now the main issue is how to reach the *island of stability* ? Because of the limitation of the nucleosynthesis by neutron capture, the only possible way to reach the stability island is nuclear reaction induced by heavy ions via complete fusion reactions the so-called *cold fusion reactions* [15]. In this method, the kinetic energy of the heavy projectile is absorbed in a compound nucleus (heavy target). It is interesting to note that the superheavy elements with $Z=105-118$ are synthesized using cold-fusion reactions [16, 17]. Although, the superheavy elements $Z=93-118$ are synthesized artificially, the knowledge about their chemistry is limited to $Z < 107$ due to the shorter β -decay life time τ_β [18]. The chemistry of elements $Z=107$ and higher Z nuclei were unknown and the prospects of the studies of their properties did not look promising because of their shorter life time. However, some isotopes of these elements are recently discovered whose life time seems to be relatively large, which gives a hope for the studies of the chemistry of the superheavy elements. In this paper, our aim is to study the properties and the stability of the recently synthesized superheavy nuclei.

The stability of a nucleus mostly depends on its structure which is again a function of the combination of number of neutron and proton. For an extremely large nucleus as it is in super-heavy region, the shape of the ground state configuration most unlikely spherical. Because of the large Coulombic repulsion, the central portion of the nucleus may take a depletion and create a bubble like structure. There are some interesting and extensive calculations for various properties including bubbles or semi-bubbles structure of superheavy elements [19, 20]. In this paper, the stability of the superheavy nuclei are discussed by two well known self-consistent mean field models, namely Skyrme Hartree-Fock (SHF) using SkI4 parameters [21] with BCS pairing and Relativistic Mean Field (RMF) model with NL3* parameter set [22].

The paper starts with a short introduction in Sec. I. The formalisms of Skyrme Hartree-Fock and relativistic mean field theory are presented in Sec.II. Mostly, the stability and structure of the synthesized nuclei will be discussed in the Results and discussions are given in Sec. III. In this section, mostly the stability and structure of the synthesized nuclei will be discussed. The bubble properties will also be analyzed taking into consideration the density distributions of proton and neutron. Sec. IV contains a brief summary of the paper.

II. THEORETICAL FORMALISMS

A. Skyrme Hartree-Fock (SHF) method

There are many known parametrization of Skyrme interaction which reproduce the experimental data for ground state properties of finite nuclei [23, 24] as well as the properties of infinite nuclear matter upto high density [25]. The general form of the Skyrme effective interaction can be expressed as a density functional $\mathcal{H}$ with some empirical parameters [24, 26, 27]:

$$\mathcal{H} = \mathcal{K} + \mathcal{H}_0 + \mathcal{H}_3 + \mathcal{H}_{eff} + \cdots, \quad (1)$$

where $\mathcal{K}$ is the kinetic energy, $\mathcal{H}_0$ the zero range, $\mathcal{H}_3$ the density dependent and $\mathcal{H}_{eff}$ the effective-mass dependent terms, which are relevant for calculating the properties of nuclear matter. More details can be found in Refs. [24, 26, 27]. These are functions of 9 parameters t_i, x_i ($i = 0, 1, 2, 3$)

and η are given as

$$\mathcal{H}_0 = \frac{1}{4}t_0 [(2+x_0)\rho^2 - (2x_0+1)(\rho_p^2 + \rho_n^2)], \quad (2)$$

$$\mathcal{H}_3 = \frac{1}{24}t_3\rho^\eta [(2+x_3)\rho^2 - (2x_3+1)(\rho_p^2 + \rho_n^2)], \quad (3)$$

$$\begin{aligned} \mathcal{H}_{eff} = & \frac{1}{8} [t_1(2+x_1) + t_2(2+x_2)] \tau \rho \\ & + \frac{1}{8} [t_2(2x_2+1) - t_1(2x_1+1)] (\tau_p \rho_p + \tau_n \rho_n). \end{aligned} \quad (4)$$

The kinetic energy $\mathcal{K} = \frac{\hbar^2}{2m}\tau$, a form used in the Fermi gas model for non-interacting Fermions. The other terms $\mathcal{H}_{S\rho}$ and $\mathcal{H}_{S\vec{J}}$ with b_4 and b'_4 are the surface contributions which also responsible for a better spin-orbit interaction are defined as

$$\begin{aligned} \mathcal{H}_{S\rho} = & \frac{1}{16} \left[3t_1(1 + \frac{1}{2}x_1) - t_2(1 + \frac{1}{2}x_2) \right] (\vec{\nabla}\rho)^2 \\ & - \frac{1}{16} \left[3t_1(x_1 + \frac{1}{2}) + t_2(x_2 + \frac{1}{2}) \right] \\ & \times \left[(\vec{\nabla}\rho_n)^2 + (\vec{\nabla}\rho_p)^2 \right], \text{ and} \end{aligned} \quad (5)$$

$$\mathcal{H}_{S\vec{J}} = -\frac{1}{2} \left[b_4\rho\vec{\nabla} \cdot \vec{J} + b'_4(\rho_n\vec{\nabla} \cdot \vec{J}_n + \rho_p\vec{\nabla} \cdot \vec{J}_p) \right]. \quad (6)$$

Here, the total nucleon number density $\rho = \rho_n + \rho_p$, the kinetic energy density $\tau = \tau_n + \tau_p$, and the spin-orbit density $\vec{J} = \vec{J}_n + \vec{J}_p$. The subscripts n and p refer to neutron and proton, respectively, and M is the nucleon mass. The $\vec{J}_q = 0$, $q = n$ or p , for spin-saturated nuclei, i.e., for nuclei with major oscillator shells completely filled. The total binding energy (BE) of a nucleus is the integral of the density functional $\mathcal{H}$. In our calculations, we have used the Skyrme Ski4 parameter set [21]. This set is known to well account the spin-orbit interaction in finite nuclei, related to the isotope shifts in Pb region and is better suited for the study of exotic nuclei. Several more recent Skyrme parameters such as SLy1-10, SkX, Ski5 and Ski6 are obtained by fitting the Hartree-Fock (HF) results with experimental data for nuclei starting from the valley of stability to neutron and proton drip-lines [21, 27–29].

B. Relativistic mean field (RMF) formalism

From last few decades, the RMF theory is applied successfully to study the structural properties of nuclei throughout the periodic table [30–33] starting from proton to neutron drip-lines. The

starting point of the RMF theory is the basic Lagrangian containing nucleons interacting with σ -, ω - and ρ -meson fields. The photon field A_μ is included to take care of the Coulomb interaction of protons. The relativistic mean field Lagrangian density is expressed as [30–34],

$$\begin{aligned}\mathcal{L} = & \bar{\psi}_i \{i\gamma^\mu \partial_\mu - M\} \psi_i + \frac{1}{2} \partial^\mu \sigma \partial_\mu \sigma - \frac{1}{2} m_\sigma^2 \sigma^2 - \frac{1}{3} g_2 \sigma^3 \\ & - \frac{1}{4} g_3 \sigma^4 - g_s \bar{\psi}_i \psi_i \sigma - \frac{1}{4} \Omega^{\mu\nu} \Omega_{\mu\nu} + \frac{1}{2} m_\omega^2 V^\mu V_\mu \\ & - g_\omega \bar{\psi}_i \gamma^\mu \psi_i V_\mu - \frac{1}{4} \vec{B}^{\mu\nu} \vec{B}_{\mu\nu} + \frac{1}{2} m_\rho^2 \vec{R}^\mu \vec{R}_\mu - \frac{1}{4} F^{\mu\nu} F_{\mu\nu} \\ & - g_\rho \bar{\psi}_i \gamma^\mu \vec{\tau} \psi_i \vec{R}^\mu - e \bar{\psi}_i \gamma^\mu \frac{(1 - \tau_{3i})}{2} \psi_i A_\mu.\end{aligned}\quad (7)$$

Here M , m_σ , m_ω and m_ρ are the masses for nucleon, σ -, ω - and ρ -mesons and ψ is its Dirac spinor. The field for the σ -meson is denoted by σ , ω -meson by V_μ and ρ -meson by R_μ . g_s , g_ω , g_ρ and $e^2/4\pi=1/137$ are the coupling constants for the σ , ω , ρ -mesons and photon respectively. g_2 and g_3 are the self-interaction coupling constants for σ mesons. By using the classical variational principle we obtain the field equations for the nucleons and mesons.

$$\begin{aligned}\{-\Delta + m_\sigma^2\} \sigma^0(r_\perp, z) = & -g_s \rho_s(r_\perp, z) \\ & - g_2 \sigma^2(r_\perp, z) - g_3 \sigma^3(r_\perp, z),\end{aligned}\quad (8)$$

$$\{-\Delta + m_\omega^2\} V^0(r_\perp, z) = g_\omega \rho_v(r_\perp, z), \quad (9)$$

$$\{-\Delta + m_\rho^2\} R^0(r_\perp, z) = g_\rho \rho_3(r_\perp, z), \quad (10)$$

$$-\Delta A^0(r_\perp, z) = e \rho_c(r_\perp, z). \quad (11)$$

The Dirac equation for the nucleons is written by

$$\{-i\alpha \cdot \nabla + V(r_\perp, z) + \beta M^\dagger\} \psi_i = \epsilon_i \psi_i. \quad (12)$$

The effective mass of the nucleon is

$$M^\dagger = M + S(r_\perp, z) = M + g_\sigma \sigma(r_\perp, z), \quad (13)$$

and the vector potential is

$$V(r_\perp, z) = g_\omega V^0(r_\perp, z) + g_\rho \tau_3 R^0(r_\perp, z) + e \frac{(1 - \tau_3)}{2} A^0(r_\perp, z). \quad (14)$$

A static solution is obtained from the equations of motion to describe the ground state properties of nuclei. The set of nonlinear coupled equations are solved self-consistently in an axially deformed

harmonic oscillator basis $N_F = N_B = 20$. The quadrupole deformation parameter β_2 is extracted from the calculated quadrupole moments of neutrons and protons through

$$Q = Q_n + Q_p = \sqrt{\frac{16\pi}{5}} \left(\frac{3}{4\pi} AR^2 \beta_2 \right), \quad (15)$$

where $R = 1.2A^{1/3}$.

The total energy of the system is given by

$$E_{total} = E_{part} + E_{\sigma} + E_{\omega} + E_{\rho} + E_c + E_{pair} + E_{c.m.}, \quad (16)$$

where E_{part} is the sum of the single particle energies of the nucleons and E_{σ} , E_{ω} , E_{ρ} , E_c , E_{pair} , E_{cm} are the contributions of the meson fields, the Coulomb field, pairing energy and the center-of-mass energy, respectively. We use the recently reported NL3* parameter set [22] in our calculations for RMF formalism [22].

III. RESULTS AND DISCUSSIONS

In the present paper, we calculate the binding energy, root mean square (rms) charge radius r_{ch} and quadrupole deformation parameter β_2 . A special attention is given to the neutron and proton density distributions of these superheavy nuclei, where we search for the bubble structure at various excited intrinsic states. The results are explained in subsection III.A to III.E.

A. Stability of nuclei

While analyzing the nuclear landscape, the stability of nucleus is maximum with a neutron to proton ratio N/Z at about 1 for light mass region. This value of N/Z increases with mass number A to neutralize the Coulomb repulsion and saturated at ~ 1.5 . For example, the neutron to proton ratio for ^{208}Pb is 1.537 and for $^{235,238}\text{U}$ these values are 1.554 and 1.587 respectively. This saturation nature of $N/Z \sim 1.5$ suggests the formation of superheavy nuclei with an $N : Z = 1.5$. To see the stability of the synthesized isotopes in the superheavy region, we plot the N/Z values in Figure 1 for $Z=105$ to 118 in addition to the predicted double close nuclei $^{292,293,304}120$. To our surprise, we find the minimum and maximum $N : Z$ as 1.433 and 1.614, respectively.

Another important quantity is the binding energy (BE) of a nucleus. This is the only experimental observable which can be measured precisely and tells many things about the stability. A




FIG. 1: The N/Z ratio of the recently synthesized nuclei with atomic number $Z = 105 - 118$. The N/Z values of the predicted superheavy isotopes $^{292,293,304}_{120}$ are also given.

nucleus is found to be more stable with a larger binding energy per particle (BE/A). The trend of BE/A as a function of mass number A is shown in Figure 2. It is clear that low mass nuclei in the valley of β -stability have more life-time compared to the heavier isotopes. This may be due to the large Coulomb repulsion of the superheavy nuclei, although the N/Z ratio is almost constant. The declining trend of BE/A indicates the difficulty in the formation of heavier superheavy nuclei. Of course, it is to be recalled that the formation of a neutron-proton bound state system requires only 2.224 MeV which is the typical binding energy of Deuteron ($BE/A \sim 1.112$ MeV) [35].

B. Shape and shape co-existence

Nuclei have different binding energy corresponding to different shape configuration in the ground or excited states. In some cases, it so happens that the binding energy of two different configurations coincide or very close to each other known as shape co-existence [38–41]. This phenomenon is more common in superheavy region, leads to complex structure of these nuclei

TABLE I: The BE (MeV), β_2 , r_{ch} (fm) (ground state) and D.F. (%) using SHF(Skl4).

Nucl.	BE	BE	BE	β_2	β_2	β_2	r_{ch}	Spherical		Prolate		Oblate		FRDM	
	Sphe.	Prolate	Oblate					D.F. _n	D.F. _p	D.F. _n	D.F. _p	D.F. _n	D.F. _p	BE	β_2
²⁶⁶ Db		1950.3	1939.7		0.262	-0.282	6.08	0.0	0.0	0.0	2.98	1.32	7.86	1953.42	0.221
²⁶⁷ Db		1956.5	1945.9		0.258	-0.283	6.09	0.0	3.41	0.0	3.41	1.29	7.43	1960.49	0.230
²⁶⁸ Db		1961.7	1951.3		0.254	-0.193	6.09	0.0	3.42	0.0	3.83	1.35	7.36	1965.86	0.221
²⁶⁹ Db		1966.9	1957.6		0.250	-0.191	6.10	0.0	3.58	0.0	3.60	1.13	7.53	1972.01	0.221
²⁷⁰ Db		1971.9	1963.7		0.259	-0.188	6.11	0.0	3.65	0.0	2.12	0.81	7.64	1976.81	0.212
²⁵⁸ Sg		1890.5	1880.0		0.287	-0.287	6.05	2.73	4.52	2.73	4.54	2.74	0.89	1895.22	0.248
²⁶⁰ Sg		1905.5	1894.9		0.284	-0.286	6.06	2.29	4.24	0.0	4.24	1.75	12.86	1909.90	0.239
²⁶² Sg		1920.1	1909.3		0.279	-0.283	6.07	1.05	3.69	1.72	3.70	1.59	12.19	1923.94	0.229
²⁷¹ Sg		1976.5	1968.0		0.254	-0.188	6.12	1.39	4.12	0.0	3.45	0.93	7.71	1981.87	0.212
²⁷⁰ Bh		1969.8	1958.7		0.254	-0.192	6.11	0.80	4.00	0.0	4.02	1.53	7.57	1973.53	0.222
²⁷² Bh		1980.8	1971.9		0.244	-0.187	6.12	0.0	4.21	0.0	4.62	1.07	7.82	1985.27	0.221
²⁷⁴ Bh		1987.4	1984.7		0.237	-0.182	6.13	0.0	4.25	0.0	3.16	0.52	8.20	1996.40	0.192
²⁶⁴ Hs		1923.6	1912.8		0.271	-0.283	6.08	0.0	4.42	1.23	3.74	1.36	12.27	1927.62	0.229
²⁶⁸ Hs		1953.4	1941.3		0.262	-0.281	6.12	0.0	3.60	0.0	4.40	1.26	11.46	1957.28	0.230
²⁷⁰ Hs		1967.4	1955.2		0.256	-0.193	6.11	0.0	4.17	0.0	4.17	1.76	7.69	1971.42	0.231
²⁷² Hs		1978.9	1968.8		0.248	-0.188	6.12	0.0	4.40	0.0	4.40	1.48	7.84	1984.20	0.222
²⁷⁵ Hs		1995.8	1988.7		0.233	-0.180	6.14	0.0	4.39	0.0	3.84	0.78	8.42	2000.78	0.183
²⁷⁴ Mt		1987.4	1978.7		0.237	-0.183	6.14	0.0	4.71	0.0	4.72	1.49	8.28	1991.77	0.222
²⁷⁵ Mt		1993.3	1985.5		0.232	-0.181	6.14	0.0	4.31	0.0	4.73	1.27	8.49	1998.36	0.212
²⁷⁶ Mt		1999.1	1992.2		0.225	-0.178	6.14	0.0	4.27	0.0	4.67	1.05	8.73	2003.64	0.202
²⁷⁸ Mt		2010.6	2005.5		0.212	-0.173	6.15	0.0	4.38	0.0	4.49	0.70	9.30	2015.32	0.136
²⁷⁰ Ds		1955.7	1945.5		0.250	-0.195	6.12	0.0	4.30	0.0	4.31	2.09	8.13	1960.53	0.221
²⁷⁹ Ds	2014.0	2011.7	2009.1	0.199	0.276	-0.171	6.20	0.0	4.29	0.0	4.81	1.08	9.70	2019.39	0.127
²⁸¹ Ds	2019.8	2021.8	2022.3	-0.030	0.340	-0.166	6.24	16.96	22.10	0.0	4.44	0.88	10.34	2031.91	0.108
²⁷⁷ Rg		1997.7	1991.3		0.220	-0.176	6.16	0.0	4.78	0.0	4.32	2.03	9.38	2003.04	0.193
²⁷⁸ Rg		2004.0	1998.4		0.220	-0.173	6.17	0.0	4.12	0.0	4.72	1.86	9.63	2008.68	0.164
²⁷⁹ Rg		2007.1	2005.4		0.288	-0.170	6.21	0.0	4.09	0.8	4.24	1.68	9.90	2015.73	0.164

Continue..

Continue of table-I

Nucl.	BE	BE	BE	β_2	β_2	β_2	$r_{ch.}$	Spherical		Prolate		Oblate		FRDM	
	Sphe.	Prolate	Oblate					D.F. _n	D.F. _p	D.F. _n	D.F. _p	D.F. _n	D.F. _p	BE	β_2
²⁸⁰ Rg		2012.5	2012.3		0.334	-0.167	6.24	0.0	4.11	0.0	4.83	1.53	10.19	2021.81	0.117
²⁸¹ Rg	2016.9	2018.4	2019.2	-0.036	0.353	-0.165	6.26	17.27	22.44	0.77	4.56	1.40	10.50	2028.85	0.099
²⁸² Rg	2023.8	2024.3	2025.9	-0.037	0.366	-0.163	6.27	17.06	20.14	0.62	4.61	1.29	10.79	2034.82	0.099
²⁸² 112	2020.4	2020.4	2022.3	-0.030	0.370	-0.161	6.28	17.85	23.20	0.0	4.75	1.94	11.07	2032.68	0.089
²⁸³ 112	2027.9	2028.2	2029.2	-0.040	0.489	-0.159	6.38	16.73	22.14	0.0	3.99	1.29	11.33	2038.67	0.089
²⁸⁴ 112	2035.1	2035.2	2036.0	-0.066	0.525	-0.159	6.41	14.18	20.42	1.41	2.78	2.34	11.93	2045.88	0.089
²⁸⁵ 112	2042.1	2041.6	2042.5	-0.068	0.532	-0.167	6.43	13.29	19.93	0.0	2.47	1.16	11.20	2051.61	0.089
²⁹⁴ 112	2099.4	2093.3	2098.9	-0.004	0.559	-0.005	6.50	3.07	15.17	0.0	5.74	0.22	11.86	2103.92	0.00
²⁸² 113	2021.0	2021.1	2017.9	-0.164	0.171	-0.159	6.18	1.02	9.57	0.0	8.43	2.71	11.44	2027.20	0.071
²⁸⁴ 113	2031.0	2034.4	2032.3	-0.002	0.158	-0.154	6.18	18.47	24.09	0.0	8.61	2.37	11.93	2040.95	0.080
²⁸⁵ 113	2038.2	2041.1	2039.2	-0.023	0.151	-0.155	6.18	18.02	23.72	0.0	8.79	2.06	11.90	2048.18	0.072
²⁸⁶ 113	2045.3	2047.7	2045.9	-0.021	0.144	-0.163	6.19	17.54	23.44	0.0	8.94	1.58	11.65	2054.22	0.072
²⁸⁶ 114	2041.4	2041.1	2042.0	-0.012	0.555	-0.150	6.46	18.59	24.53	0.0	0.0	2.63	12.55	2051.59	-0.096
²⁸⁷ 114	2048.6	2047.7	2048.8	-0.012	0.554	-0.158	6.46	18.06	24.19	0.0	4.82	2.06	12.13	2057.65	-0.078
²⁸⁸ 114	2055.8	2054.1	2055.5	-0.024	0.547	-0.165	6.46	16.76	23.18	0.0	4.63	2.14	12.09	2065.01	0.053
²⁸⁹ 114	2062.8	2060.3	2062.2	-0.011	0.537	-0.149	6.46	16.07	22.95	0.0	4.81	2.23	12.36	2071.04	-0.052
²⁹⁶ 114	2108.9	2102.3	2108.4	-0.003	0.565	-0.043	6.53	4.05	16.58	0.0	4.32	1.68	13.68	2113.22	0.00
²⁹⁸ 114	2121.1	2113.3	2120.8	0.000	0.566	-0.012	6.54	0.0	11.18	0.0	4.72	0.0	10.74	2123.30	0.00
²⁸⁷ 115	2043.1	2044.9	2044.1	-0.008	0.153	-0.137	6.20	19.86	25.48	0.0	6.57	4.18	13.69	2052.72	-0.096
²⁸⁸ 115	2050.5	2051.9	2051.2	-0.005	0.145	-0.137	6.20	19.24	25.09	0.0	7.07	3.66	13.38	2059.12	-0.087
²⁸⁹ 115	2057.8	2058.8	2057.9	-0.002	0.137	-0.141	6.21	18.19	25.44	0.0	7.48	3.02	12.92	2066.45	0.053
²⁹⁰ 115	2064.9	2064.7	2065.2	0.000	0.117	-0.119	6.20	16.88	23.61	2.34	10.75	3.83	13.59	2072.59	-0.079
²⁹¹ 115	2071.9	2071.8	2072.4	0.000	0.064	-0.102	6.20	15.46	22.70	8.62	19.14	4.39	14.44	2079.83	-0.061
²⁹⁰ 116	2059.5	2060.5	2060.3	0.002	0.134	-0.121	6.21	17.92	24.59	0.0	7.90	4.92	15.51	2068.76	0.072
²⁹¹ 116	2066.7	2066.7	2067.5	0.002	0.110	-0.108	6.21	16.61	23.78	0.3	11.81	5.56	15.29	2074.84	0.072
²⁹² 116	2073.9	2074.1	2074.7	0.002	0.062	-0.097	6.21	15.32	22.96	10.18	19.89	5.37	15.45	2082.48	-0.070
²⁹³ 116	2080.9	2081.1	2081.8	0.002	0.052	-0.092	6.21	13.55	21.98	10.76	2.35	4.07	14.79	2088.40	-0.070

Continue...




FIG. 2: The binding energy per nucleon (BE/A) is compared with the experimental data. The stars and circles are for RMF(NL3*) and SHF(SkI4), respectively.

which provides the information about the oscillation of nuclei between two or three existing shape by perturbing small energy. Here, in our studied nuclei, we find many such examples, where the ground and first excited binding energies are degenerated. Also, in several cases the nuclei have well defined three distinct solutions (oblate, spherical, prolate) with almost same energy which strongly indicates the nuclei may oscillate from oblate to spherical to prolate configuration.

For analyzing the result, we took a small binding energy difference around ≤ 2 MeV to take care of shape co-existence for our calculations. Because of this small difference in binding energies between these two solutions, the ground state can be changed to the low-lying excited state or vice-versa by a small change in input parameters like pairing gaps etc. It is seen from Tables I and II the shape co-existence is observed for $^{277,278,281,282}\text{Rg}$, $^{282,283,284,285}\text{112}$, $^{286,287,288}\text{114}$, $^{294,297}\text{118}$ and $^{292,293}\text{120}$ in SHF calculations and RMF also produces the shape co-existence for $^{294}\text{112}$, $^{286}\text{113}$, $^{288,289,296,298}\text{114}$, $^{290,291}\text{115}$, $^{291,292}\text{116}$, $^{293,294,297}\text{117}$, $^{297}\text{118}$, $^{292,293}\text{120}$. A shape co-existence of nuclei indicates that there is a competition among the different shapes to acquire the possible ground state for maximum stability and the final shape could be an admixture of these low-lying

Continue of Table- I

Nucl.	BE	BE	BE	β_2	β_2	β_2	$r_{ch.}$	Spherical		Prolate		Oblate		FRDM	
	Sphe.	Prolate	Oblate					D.F. _n	D.F. _p	D.F. _n	D.F. _p	D.F. _n	D.F. _p	BE	β_2
²⁹³ 117	2075.6	2074.3	2076.4	0.039	0.536	-0.099	6.50	13.64	22.17	0.0	0.0	4.57	14.65	2083.06	-0.087
²⁹⁴ 117	2082.8	2080.9	2083.6	0.043	0.538	-0.095	6.51	12.18	21.27	0.0	0.0	2.99	13.77	2089.22	-0.087
²⁹⁷ 117	2103.6	2100.6	2103.3	0.031	0.561	-0.083	6.55	8.27	18.36	0.0	0.0	0.0	11.86	2109.27	-0.008
²⁹⁴ 118	2076.9	2076.1	2077.7	0.055	0.548	-0.102	6.52	9.32	19.71	0.0	0.34	3.74	13.73	2084.78	-0.087
²⁹⁷ 118	2098.2	2096.6	2098.6	-0.023	0.561	-0.087	6.55	11.38	20.22	0.0	1.28	0.0	11.30	2104.59	-0.035
²⁹² 120	2046.4	2047.1	2047.3	-0.006	0.530	-0.125	6.50	26.49	31.54	0.4	0.0	7.58	15.13	2055.19	-0.130
²⁹³ 120	2054.5	2054.9	2055.2	-0.005	0.528	-0.124	6.51	25.35	30.75	0.3	0.98	6.71	14.56	2062.42	0.089
³⁰⁴ 120	2134.9	2132.5	2132.7	-0.008	0.558	-0.074	6.59	0.0	8.36	0.0	3.62	0.0	10.01	2137.99	0.00

bands.

C. Binding energy and quadrupole deformation parameter

The ground state binding energy and quadrupole deformation parameter for all the observed nuclei considered here are given in Tables I and II. The much discussed superheavy isotopes ^{292,304}120 which are considered to be the next double closed nuclei beyond ²⁰⁸Pb are also listed. To find the ground state solution, the calculations are done with an initial spherical, prolate and oblate quadrupole deformation parameter β_0 both in the Skyrme Hartree-Fock (SHF) and relativistic mean field (RMF) formalisms. The well known SkI4[21] and NL3*[22] parametrizations are used through out the SHF and RMF calculations, respectively. It is to be noted that the maximum binding energy corresponds to the ground state and all other solutions are the intrinsic excited state configurations. Considering this criteria, we evaluate the ground state binding energy and also the first and second excited states wherever available. The results obtained from SHF are listed in Table I and those for RMF are listed in Table II. Since the experimental binding energy of these nuclei are not known, to get a first hand information about the predictive power of our calculations, we have inserted the most commonly used binding energy of finite range droplet model (FRDM) [36]. The quadrupole deformation parameter β_2 obtained in FRDM is also given for comparison [36, 37]. Comparing Tables I and II, it is remarkable to note that the binding energy obtained by

TABLE II: Same as Table I with RMF(NL3*) model.

Nucleus	BE	BE	BE	β_2	β_2	β_2	$r_{ch.}$	Spherical		Prolate		Oblate	
	Spherical	Prolate	Oblate					(D.F.) _n	(D.F.) _p	(D.F.) _n	(D.F.) _p	(D.F.) _n	(D.F.) _p
²⁶⁶ Db	1941.5	1955.9	1946.2	0.002	0.273	-0.308	6.129	19.57	18.00	1.22	2.95	5.22	20.05
²⁶⁷ Db	1947.9	1961.6	1952.1	0.002	0.269	-0.307	6.133	19.79	18.11	1.04	1.91	5.28	15.67
²⁶⁸ Db	1954.3	1966.7	1957.9	0.001	0.267	-0.306	6.138	19.98	18.29	0.92	2.92	4.08	20.32
²⁶⁹ Db	1960.4	1971.4	1963.6	0.001	0.263	-0.306	6.143	20.20	18.52	0.87	1.55	4.19	20.66
²⁷⁰ Db	1966.5	1976.0	1969.1	0.002	0.256	-0.306	6.147	20.40	18.78	0.87	1.23	4.27	21.14
²⁵⁸ Sg	1883.6	1899.6	1889.5	0.004	0.291	-0.301	6.091	17.28	16.26	6.26	4.43	9.54	18.95
²⁶⁰ Sg	1898.1	1913.9	1903.9	0.002	0.285	-0.301	6.102	18.04	16.86	6.18	4.46	7.35	19.66
²⁶² Sg	1912.1	1927.7	1917.7	0.002	0.281	-0.305	6.113	18.68	17.38	5.32	4.33	5.69	19.37
²⁷¹ Sg	1971.5	1980.9	1973.6	0.002	0.254	-0.309	6.157	20.68	19.31	0.38	3.48	5.15	20.99
²⁷⁰ Bh	1963.1	1975.3	1965.9	0.002	0.262	-0.311	6.158	0.42	19.16	0.60	4.21	4.58	20.16
²⁷² Bh	1975.9	1985.4	1977.8	0.002	0.244	-0.312	6.164	20.87	19.69	0.41	4.11	4.89	20.86
²⁷⁴ Bh	1995.9	1990.8	1989.3	0.205	0.216	-0.319	6.162	1.00	6.77	0.54	0.26	5.99	22.15
²⁶⁴ Hs		1932.1	1922.6		0.269	-0.311	6.130	5.72	7.62	6.06	4.62	5.28	19.75
²⁶⁸ Hs		1959.9	1950.1		0.265	-0.315	6.152	1.98	5.76	1.86	3.08	3.64	19.94
²⁷⁰ Hs		1972.9	1963.1		0.260	-0.314	6.161	0.78	5.24	0.88	4.85	4.34	20.10
²⁷² Hs		1984.1	1975.6		0.249	-0.313	6.169	0.43	5.29	0.99	4.78	3.55	20.51
²⁷⁵ Hs		2000.6	1993.6		0.203	-0.322	6.171	1.19	7.66	1.40	8.53	5.89	22.19
²⁷⁴ Mt		1993.4	1985.1		0.216	-0.319	6.173	1.08	7.69	1.49	1.16	5.09	21.46
²⁷⁵ Mt		1999.1	1991.3		0.207	-0.324	6.176	1.22	8.00	1.47	0.96	5.69	22.16
²⁷⁶ Mt		2004.7	1997.4		0.199	-0.327	6.180	1.47	8.16	1.43	8.98	6.16	22.77
²⁷⁸ Mt	2015.8	2013.6	2009.4	0.185	0.373	-0.330	6.186	1.39	8.30	1.06	3.68	6.53	23.77
²⁷⁰ Ds		1963.9	1954.9		0.251	-0.319	6.167	2.84	7.77	2.88	8.09	4.35	20.93
²⁷⁹ Ds	2020.0	2017.4	2013.3	0.182	0.384	-0.336	6.196	1.40	8.57	2.63	3.88	6.81	24.27
²⁸¹ Ds	2031.2	2029.2	2025.4	0.168	0.476	-0.334	6.202	1.65	8.86	4.93	7.06	6.76	24.47
²⁷⁷ Rg	2005.6	2002.7	1997.8	0.199	0.341	-0.345	6.194	1.39	8.61	2.16	6.57	7.15	25.13
²⁷⁸ Rg	2011.8	2008.8	2004.2	0.191	0.349	-0.345	6.198	1.41	8.69	2.23	6.55	7.18	25.22
²⁷⁹ Rg	2017.9	2014.7	2010.7	0.185	0.362	-0.345	6.202	1.41	8.73	2.38	6.58	7.18	25.26

Continue...

Continue of Table II

Nucleus	BE	BE	BE	β_2	β_2	β_2	$r_{ch.}$	Spherical		Prolate		Oblate	
	Spherical	Prolate	Oblate					(D.F.) _n	(D.F.) _p	(D.F.) _n	(D.F.) _p	(D.F.) _n	(D.F.) _p
²⁸⁰ Rg	2023.9	2020.6	2017.0	0.179	0.385	-0.343	6.206	1.41	8.79	2.64	6.42	7.16	25.27
²⁸¹ Rg	2029.7	2027.2	2023.3	0.173	0.503	-0.341	6.209	1.46	8.89	6.42	8.57	7.09	25.22
²⁸² Rg	2035.4	2033.3	2029.4	0.166	0.497	-0.339	6.212	1.59	9.04	5.86	8.17	7.05	25.23
²⁸² 112	2033.3	2030.8	2026.8	0.171	0.515	-0.346	6.219	1.45	9.06	6.79	8.79	7.33	25.95
²⁸³ 112	2039.2	2037.2	2033.2	0.165	0.508	-0.343	6.222	1.53	9.17	6.22	7.65	7.29	25.92
²⁸⁴ 112	2041.3	2043.4	2039.3	0.002	0.501	-0.340	6.436	23.96	22.74	5.72	8.33	7.25	25.95
²⁸⁵ 112	2047.4	2049.5	2045.3	0.002	0.496	-0.337	6.438	23.56	22.59	5.38	8.38	7.15	26.04
²⁹⁴ 112	2097.6	2097.3	2098.6	0.001	0.516	-0.024	6.240	8.94	17.06	6.11	9.59	7.89	17.18
²⁸² 113	2030.3	2025.4	2023.6	0.175	0.363	-0.352	6.226	1.42	9.04	2.27	7.30	7.45	26.41
²⁸⁴ 113	2041.9	2042.5	2036.5	0.139	0.164	-0.347	6.233	4.71	11.34	1.55	1.03	7.46	26.42
²⁸⁵ 113	2045.1	2048.5	2042.8	0.001	0.158	-0.344	6.236	24.18	22.89	2.05	10.44	7.46	26.48
²⁸⁶ 113	2051.4	2053.4	2051.5	0.002	0.502	-0.148	6.454	24.18	22.88	5.34	8.18	9.97	14.25
²⁸⁶ 114	2048.6	2051.6	2046.0	0.001	0.158	-0.347	6.246	25.21	23.32	1.73	10.37	7.57	26.82
²⁸⁷ 114	2054.9	2057.6	2054.8	0.001	0.151	-0.173	6.249	24.84	23.22	2.39	10.56	8.18	17.80
²⁸⁸ 114	2061.1	2063.1	2061.3	0.002	0.506	-0.184	6.475	24.10	22.91	4.86	5.12	7.17	14.84
²⁸⁹ 114	2067.1	2069.1	2067.3	0.002	0.508	-0.191	6.482	23.24	22.60	4.66	7.25	6.56	14.98
²⁹⁶ 114	2107.5	2107.5	2105.1	0.000	0.538	-0.206	6.257	10.22	18.10	5.37	7.44	3.08	15.19
²⁹⁸ 114	2118.0	2117.9	2114.5	0.000	0.573	-0.316	6.265	1.31	12.94	5.53	8.09	9.02	29.45
²⁸⁷ 115	2051.7	2053.8	2051.5	0.001	0.159	-0.104	6.257	25.91	23.67	1.01	9.99	17.34	18.44
²⁸⁸ 115	2058.2	2060.1	2057.9	0.001	0.152	-0.180	6.259	25.52	23.54	1.81	10.25	7.98	15.41
²⁸⁹ 115	2064.5	2066.1	2064.5	0.002	0.143	-0.187	6.261	24.84	23.31	2.69	10.65	7.24	15.47
²⁹⁰ 115	2070.6	2072.4	2070.7	0.002	0.515	-0.195	6.499	23.96	22.98	4.05	3.20	6.64	15.61
²⁹¹ 115	2077.3	2078.5	2076.8	0.073	0.518	-0.213	6.507	13.28	18.94	3.92	5.22	6.39	16.10
²⁹⁰ 116	2067.5	2068.1	2067.2	0.012	0.144	-0.187	6.271	25.22	23.65	2.20	10.26	7.53	16.11
²⁹¹ 116	2073.9	2075.3	2073.7	0.044	0.522	-0.194	6.561	20.61	22.01	3.40	3.65	6.92	16.17
²⁹² 116	2080.3	2081.7	2079.9	0.052	0.525	-0.212	6.524	17.45	19.62	3.31	3.41	6.82	16.53
²⁹³ 116	2086.6	2087.8	2086.2	0.053	0.529	-0.225	6.532	16.60	20.59	3.32	3.29	6.81	16.72

Continue...




FIG. 3: The neutron, proton and total matter density distribution for some of the selected superheavy nuclei using SHF(SkI4).

SHF(SkI4) and RMF(NL3*) are in excellent agreement with FRDM prediction. The deviation of our results (from the FRDM prediction) is a maximum of about 0.5%. A further inspection of the results finds an edge of RMF(NL3*) over SHF(SkI4). In general, the calculated binding energies and deformation parameters are almost agree with each other in all the three models.

D. Density and bubble structure

The density of a nucleus has the gross information about the size, shape and distribution of nucleons. The density distribution from SHF(SkI4) and RMF(NL3*) for some of the selected nuclei are given in Figures 3 – 7 obtained for all the three solutions, i.e. spherical, prolate and oblate. Normally, the density at the center of nucleus has the maximum value and decays to zero at the surface. However, in some specific cases, e.g. for spherical solutions of $^{285}_{113}$, $^{294}_{117}$, $^{292,293}_{120}$, this trend of density distribution shows an anomalous behavior, i.e. a dip at the center and a hump

Continue of Table II

Nucleus	BE	BE	BE	β_2	β_2	β_2	$r_{ch.}$	Spherical		Prolate		Oblate	
	Spherical	Prolate	Oblate					(D.F.) _n	(D.F.) _p	(D.F.) _n	(D.F.) _p	(D.F.) _n	(D.F.) _p
²⁹³ 117	2082.8	2084.4	2080.2	0.000	0.531	-0.355	6.540	24.50	23.45	2.95	2.06	7.83	28.12
²⁹⁴ 117	2089.4	2090.7	2086.6	0.043	0.535	-0.353	6.550	19.07	21.64	3.00	2.40	7.77	28.33
²⁹⁷ 117	2107.5	2108.9	2105.3	0.006	0.550	-0.352	6.580	19.05	21.84	3.75	3.33	7.62	28.81
²⁹⁴ 118	2106.2	2095.8	2094.2	0.001	0.529	-0.235	6.532	15.06	20.13	4.20	5.02	5.79	15.99
²⁹⁷ 118	2111.9	2112.2	2109.2	0.000	0.567	-0.355	6.582	10.88	18.38	4.78	5.64	4.16	15.53
²⁹² 120	2061.5	2060.8	2057.9	0.000	0.547	-0.392	6.281	28.92	26.53	3.49	1.47	3.51	21.78
²⁹³ 120	2068.7	2068.3	2065.1	0.000	0.542	-0.388	6.285	28.56	26.49	3.09	1.48	3.75	29.11
³⁰⁴ 120	2139.3	2142.4	2137.4	0.000	0.592	-0.369	6.678	3.68	16.00	3.44	0.81	8.33	29.69

nearby to it following a slow decreasing in density to zero at the surface. This type of density distribution is known as bubble structure [42]. The first possibility for existence of toroidal or bubble structure other than the spherical topological structure was suggested by Wheeler [42]. The occurrence of bubble nuclei has been extensively studied by Wilson [43] and later by Siemens and Bethe [44]. Several models like independent particle model [45] and Hartree Fock Model [46] also investigated the possibility of low density region at the center of the nuclei. This structure is not confined to a particular region, but have the possibility for light to superheavy nuclei [47, 48]. The contribution of density at $r = 0$ is offered by non zero wavefunction of s-states and depopulation of this level leads to the depletion of central density and formed a bubble like structure [48]. The inversion of s-states to other higher states may be the possibility of depletion of central density and can be interpreted as *s-d* orbital inversion [49, 50].

The bubble effect can be realized quantitatively by calculating depletion fraction for the neutrons as well as protons using the relation [48]

$$(D.F.)_{\alpha} = \frac{(\rho_{max})_{\alpha} - (\rho_{cen})_{\alpha}}{(\rho_{max})_{\alpha}} \times 100, \quad (17)$$

where ρ_{max} , ρ_{cen} represent the maximum and central density respectively and α denotes neutron or proton. The calculated values of depletion fraction for neutron as well as proton by using non-relativistic and relativistic densities are given in Tables I and II for all three solutions. In our present study, most of the nuclei achieve the prolate shape as their ground state solution both in the SHF and RMF calculations. Here, we do not find significant bubble structure, i.e. no good




FIG. 4: Same as Fig. 3, but for $^{292,293,304}_{120}$ with SHF(SkI4).

amount of depletion fraction are estimated for this shape, except a few cases. It is evident from Tables I and II, a remarkable depletion fraction is there in $^{292,293}_{120}$ over the whole series of nuclei on spherical solution in both models.

E. Q_α and T_α for superheavy nuclei

The Q_α value of a nucleus provides valuable information for its stability. This is more important for superheavy nuclei, which is the central theme of study in the present paper. Thus, it is worthwhile to evaluate the Q_α values for the nuclei which will also give us to estimate the life time T_α of the considered nuclei. It is to be noted that α -decay is the decay mode for these elements (and not the β -decay) which lie on the valley of stability with a neutron to proton ratio ~ 1.5 (See Fig. 1). The calculated Q_α and T_α for $Z = 105 - 120$ within the relativistic and non-relativistic formalism using NL3* and SkI4 parameters are listed in Table III. The Q_α is calculated by using the following expression [51]

$$Q_\alpha = BE(N, Z) - BE(N - 2, Z - 2) - BE(2, 2). \quad (18)$$

TABLE III: The calculated Q_α and life time T_α using SHF(SkI4) and RMF(NL3*) are compared with FRDM and experimental data.

Nucleus	Q_α (SHF)	Q_α (RMF)	Q_α (FRDM)	Q_α (exp.)	T_α (SHF)	T_α (RMF)	T_α (FRDM)	T_α (exp.)
²⁶⁶ Db	7.64	7.17	7.36		1.03×10^6 s	8.87×10^7 s	$10^{7.15}$ s	
²⁶⁷ Db	7.41	7.15	7.09		3.94×10^6 s	4.84×10^7 s	$10^{7.96}$ s	
²⁶⁸ Db	8.01	7.49	7.19		40673.0s	4.23×10^6 s	$10^{7.87}$ s	
²⁶⁹ Db	8.41	7.98	7.59		717.39s	2.32×10^4 s	$10^{6.95}$ s	
²⁷⁰ Db	8.38	8.19	7.84	8.80 ± 0.1	1995.7s	9073.49s	$10^{5.25}$ s	1.3min
²⁵⁸ Sg	10.02	9.48	9.45		4.05ms	0.13s	$10^{-0.83}$ s	
²⁶⁰ Sg	9.49	9.33	9.93		0.12s	0.34s	$10^{-2.15}$ s	
²⁶² Sg	8.80	9.04	9.61		14.42s	2.63s	$10^{-1.26}$ s	
²⁷¹ Sg	8.84	8.88	8.40	8.54 ± 0.08	125.06s	95.39s	$10^{3.55}$ s	$1.9^{+2.4}_{-0.6}$ s
²⁷⁰ Bh	8.80	8.92	8.18	8.93 ± 0.08	427.56s	173.58s	$10^{4.76}$ s	87.98s
²⁷² Bh	9.20	9.61	8.88	9.01	23.83s	1.49s	$10^{2.36}$ s	11.8s
²⁷⁴ Bh	8.60	8.34	8.71	8.80 ± 0.1	1953.31s	14779.07s	$10^{2.93}$ s	1.3min
²⁶⁴ Hs	10.20	10.06	10.57		5.94ms	14.3ms	$10^{-3.18}$ s	
²⁶⁸ Hs	9.12	9.59	9.00		7.16s	28.2ms	$10^{1.24}$ s	
²⁷⁰ Hs	8.58	9.44	8.69		406.49s	79.2ms	$10^{2.23}$ s	
²⁷² Hs	9.98	10.08	9.20		22.95ms	12.1ms	$10^{0.61}$ s	
²⁷⁵ Hs	9.00	8.75	9.39	9.30 ± 0.06	198.15s	1255.21s	$10^{1.09}$ s	$0.19^{+0.22}_{-0.07}$ s
²⁷⁴ Mt	10.70	10.13	10.06	10.02 ± 1.08	8.48ms	249.99ms	$10^{-0.41}$ s	472.6ms
²⁷⁵ Mt	10.34	9.60	10.06	9.67	0.03 s	3.30s	$10^{-0.77}$ s	0.98s
²⁷⁶ Mt	10.00	8.92	9.93	9.71	0.55s	906.38s	$10^{-0.05}$ s	0.72s
²⁷⁸ Mt	9.30	8.45	9.37	9.55 ± 0.19	57.68s	34927.55s	$10^{1.54}$ s	11.0s
²⁷⁰ Ds	11.38	10.41	10.31		3.19×10^5 s	7.1ms	$10^{-1.89}$ s	
²⁷⁹ Ds	10.10	8.85	9.68	9.70 ± 0.06	55.67ms	3047.07s	$10^{0.91}$ s	$0.2^{+0.05}_{-0.04}$ s
²⁸¹ Ds	13.34	8.37	8.55	8.83	4.21×10^{-8} s	1.39×10^5 s	$10^{4.52}$ s	1.6min
²⁷⁷ Rg	12.04	10.04	11.50		1.31×10^{-5} s	0.865s	$10^{-3.72}$ s	
²⁷⁸ Rg	11.70	9.89	11.39	10.69 ± 0.08	0.15s	4.93s	$10^{3.11}$ s	6.2ms
²⁷⁹ Rg	11.30	9.52	10.92	10.37	0.56s	27.53s	$10^{-2.36}$ s	0.17s

Continue...

Continue of Table- III

Nucleus	Q_α (SHF)	Q_α (RMF)	Q_α (FRDM)	Q_α (exp.)	T_α (SHF)	T_α (RMF)	T_α (FRDM)	T_α (exp.)
^{280}Rg	10.80	9.20	10.13	9.75	19.59s	557.61s	$10^{0.03}\text{s}$	5.1s
^{281}Rg	14.77	12.99	9.37		$1.55 \times 10^{-10}\text{s}$	$1.64 \times 10^{-7}\text{s}$	$10^{1.90}\text{s}$	
^{282}Rg	14.60	8.76	8.79	9.00 ± 0.1	$6.29 \times 10^{-10}\text{s}$	15526.06s	$10^{4.10}\text{s}$	0.74s
$^{282}_{112}$	15.87	9.37	9.42		$1.04 \times 10^{-12}\text{s}$	29.46s	$10^{1.30}\text{s}$	
$^{283}_{112}$	14.10	9.09	9.01	9.34 ± 0.06	$6.32 \times 10^{-9}\text{s}$	2666.54s	$10^{3.66}\text{s}$	$3.8^{+1.2}_{-0.7}\text{s}$
$^{284}_{112}$	11.48	10.57	8.69	9.17 ± 0.05	$7.15 \times 10^{-5}\text{s}$	0.012s	$10^{3.67}\text{s}$	$9.8^{+18}_{-3.8}\text{s}$
$^{285}_{112}$	8.00	9.98	8.59	9.15	$1.97 \times 10^7\text{s}$	5.44s	$10^{5.09}\text{s}$	29s
$^{294}_{112}$	6.12	7.71	7.53		$1.69 \times 10^{15}\text{s}$	$2.46 \times 10^7\text{s}$	$10^{8.15}\text{s}$	
$^{282}_{113}$	11.12	9.89	9.78	10.62 ± 0.08	8.20ms	21.28s	$10^{1.66}\text{s}$	88.9ms
$^{284}_{113}$	10.50	9.68	9.15	9.97 ± 0.05	46.84ms	94.43s	$10^{3.61}\text{s}$	1.36s
$^{285}_{113}$	5.60	9.45	8.97	9.48 ± 0.11	$5.74 \times 10^{19}\text{s}$	203.59s	$10^{3.86}\text{s}$	7.9s
$^{286}_{113}$	4.00	10.34	8.90	9.63 ± 0.1	$9.55 \times 10^{33}\text{s}$	1.27s	$10^{4.44}\text{s}$	28.3s
$^{286}_{114}$	7.60	10.05	9.39	10.19 ± 0.06	$4.57 \times 10^8\text{s}$	1.26s	$10^{2.08}\text{s}$	$0.13^{+0.01}_{-0.01}\text{s}$
$^{287}_{114}$	8.80	9.94	9.31	9.54	$1.23 \times 10^5\text{s}$	29.91s	$10^{3.39}\text{s}$	3.8s
$^{288}_{114}$	7.70	8.61	9.16	9.84 ± 0.05	$1.71 \times 10^8\text{s}$	$4.83 \times 10^4\text{s}$	$10^{2.80}\text{s}$	$1.9^{+3.8}_{-0.8}\text{s}$
$^{289}_{114}$	7.60	8.66	8.87	9.82	$5.33 \times 10^9\text{s}$	$3.67 \times 10^5\text{s}$	$10^{4.86}\text{s}$	2.6s
$^{296}_{114}$	6.75	8.32	8.68		$4.68 \times 10^{12}\text{s}$	$5.864 \times 10^5\text{s}$	$10^{4.43}\text{s}$	
$^{298}_{114}$	15.6	7.91	8.91		$6.92 \times 10^{-12}\text{s}$	$2.389 \times 10^7\text{s}$	$10^{3.63}\text{s}$	
$^{287}_{115}$	7.09	11.01	10.25	10.59	$1.49 \times 10^{12}\text{s}$	42.8ms	$10^{0.63}\text{s}$	32ms
$^{288}_{115}$	10.80	10.75	10.12	10.48	318.88ms	426.6ms	$10^{1.33}\text{s}$	250ms
$^{289}_{115}$	10.60	10.74	10.03	10.31 ± 0.09	482.31ms	204.84ms	$10^{1.26}\text{s}$	320ms
$^{290}_{115}$	12.20	9.23	9.92	9.95 ± 0.4	0.165s	11050.16s	$10^{1.90}\text{s}$	0.023s
$^{291}_{115}$	8.86	10.15	9.66		87783.45s	8.45s	$10^{2.34}\text{s}$	
$^{290}_{116}$	8.90	11.76	11.12	10.84 ± 0.08	24443.12s	0.23ms	$10^{-2.12}\text{s}$	$7.1^{+3.2}_{-1.7}\text{s}$
$^{291}_{116}$	8.50	10.58	11.11	10.74 ± 0.07	$7.77 \times 10^6\text{s}$	2.25s	$10^{-1.03}\text{s}$	18^{+22}_{-6}ms
$^{292}_{116}$	9.40	9.689	10.82	10.71 ± 0.15	528.65s	66.091s	$10^{-1.37}\text{s}$	33^{+155}_{-15}ms
$^{293}_{116}$	9.30	9.679	10.94	10.67 ± 0.06	12923.99s	825.50s	$10^{-0.59}\text{s}$	53^{+62}_{-19}ms

Continue...




FIG. 5: Same as Fig. 3, but with RMF(NL3*).

Continue of Table- III

Nucleus	Q_α (SHF)	Q_α (RMF)	Q_α (FRDM)	Q_α (exp.)	T_α (SHF)	T_α (RMF)	T_α (FRDM)	T_α (exp.)
$^{293}_{117}$	10.70	9.989	11.68	11.03 ± 0.08	1.07s	104.71s	$10^{-2.40}$ s	21ms
$^{294}_{117}$	9.40	10.00	11.67	10.81 ± 0.1	15027.71s	213.62s	$10^{-2.05}$ s	112ms
$^{297}_{117}$	10.74	9.52	11.74		0.844s	2869.78s	$10^{-2.55}$ s	
$^{294}_{118}$	11.10	9.60	12.28	10.84	33.36ms	553.44s	$10^{-4.24}$ s	7.1ms
$^{297}_{118}$	4.40	9.95	12.10		5.13×10^{32} s	555.59s	$10^{-2.79}$ s	
$^{292}_{120}$	12.42	10.37	13.89		9.68×10^{-5} s	12.30s	$10^{-6.96}$ s	
$^{293}_{120}$	12.24	11.05	13.69		2.81ms	2.09s	$10^{-5.51}$ s	
$^{304}_{120}$	12.07	10.19	13.82		0.58ms	42.53s	$10^{-6.83}$ s	

Here, $BE(N,Z)$ and $BE(N-2,Z-2)$ are the binding energy of the parent and daughter nuclei, respectively. $BE(2,2)$ is the binding energy of α particle (^4He) i.e. 28.3 MeV [35]. We are using the




FIG. 6: Same as Fig. 3, but for $^{292,293,304}_{120}$ with RMF(NL3*).

empirical Viola-Seaborg formula [52, 53] for the calculation of T_α which is define as:

$$\log_{10}(T_\alpha) = \frac{(aZ + b)}{\sqrt{Q_\alpha}} + (cZ + d) + h \log, \quad (19)$$

where Z is the charge number of the parent nucleus and a, b, c, d are the fitting constants defined as [52–54] $a = 1.66175, b = -8.5166, c = -0.20228, d = -33.9069$. The last term in above equation is known as even-odd hindrance factor and is obtained by fits to odd nuclei. It takes the value as $h \log = 0.0$ (for $Z = \text{even}, N = \text{even}$), $h \log = 0.772$ (for $Z = \text{odd}, N = \text{even}$), $h \log = 1.066$ (for $Z = \text{even}, N = \text{odd}$), $h \log = 1.114$ (for $Z = \text{odd}, N = \text{odd}$). From the above equation, one can find that $T_\alpha \propto 10^{\frac{1}{\sqrt{Q_\alpha}}}$, hence a small change in Q_α creates a large difference in T_α . This observation reflects in our results (see Table III).

IV. SUMMARY

In summary, we study a broad spectrum of the superheavy nuclei for $Z = 105 - 120$ in the frame work of non-relativistic and relativistic mean field formalisms. The calculations are done in an axially deformed coordinates using constant gap BCS pairing. Except $Z=120$, all other isotopes




FIG. 7: The neutron, proton and total matter density distribution for $^{293}_{117}$ with RMF(NL3*).

are synthesized artificially whose properties are studied. From the calculated binding energies, Q_α and T_α are estimated. The results are compared with the most acceptable FRDM data as well as with the experimental observations, wherever available. Over all comparison of the results are quite agreeable with each other in all the three models along with the experimental data. Another important observation of our present analysis is the constant value of neutron to proton ratio of ~ 1.5 for all the observed nuclei. This is enough indication to conclude that the superheavy nuclei which will be synthesized in future will have neutron to proton ratio of the order of 1.5. However, the binding energy per particle will have much less than the present finding of $BE \sim 7$ MeV.

On the basis of the depletion of central density, a bubble structure is studied for some superheavy nuclei in the excited states. This observation is model independent. The maximum depletion fraction calculated in the considered nuclei is for $^{292,293}_{120}$. In some of the cases, [Fig. 7] a depletion of proton density nearer to the center (but not at the center exactly) indicates a special type of proton distribution. In this case, the center is slightly bulgy and a considerably depression afterward. Again a big hump almost at the mid distance of the center and the surface appears in the density distribution for the spherical state. This feature of the nucleus is unique and need more

investigation. Work in this direction is in progress [55].

V. ACKNOWLEDGMENTS

One of the author (MI) acknowledge the hospitality provided by Institute of Physics, Bhubaneswar during the work.

-
- [1] Scharff-Goldhaber, *Nucleonics* **15**, 122 (1957).
 - [2] S. G. Nilsson, C. F. Tsang, A. Sobiczewski, Z. Szymanski, S. Wycech, C. Gustafson, I. -L. Lamm, P. Möller and B. Nilsson, *Nucl. Phys. A* **131**, 1 (1969).
 - [3] G. Herrmann, *Nature* **280**, 543 (1979).
 - [4] K. Kumar, *Superheavy Elements* (IOP Publishing Ltd. 1989); *Physica Scripta*, **T32**, 31 (1990); S. Hofmann, *Rus. Chem. Rev.* **78**, 1123 (2009).
 - [5] W. D. Myers and W. J. Swiatecki, *Nucl. Phys.* **81**, 1 (1966).
 - [6] H. W. Meldner, *Ark. Fys.* **36**, 593 (1978).
 - [7] H. W. Meldner, in *Superheavy Elements* ed. M. A. K. Lodhi (Oxford: Pergamon) p. 495.
 - [8] H. S. Köhler, *Nucl. Phys. A* **162**, 385 (1971).
 - [9] S. G. Nilsson, *Mat. Fys. Medd. Dan. Vid. Selsk* **29**, 16 (1955).
 - [10] B. R. Mottelson and S. G. Nilsson, *Phys. Rep.* **99**, 1615 (1955); *Mat. Fys. Skr. Dan. Vid. Selsk*, **8**, 1 (1959).
 - [11] G. Andersson, S. E. Larsson, G. Leander, S. G. Nilsson, I. Ragnarsson and S. Aberg, *Phys. Lett. B* **65**, 209 (1976).
 - [12] V. M. Strutinsky, *Nucl. Phys. A* **95**, 420 (1967); **122**, 1 (1968).
 - [13] M. Bhunya and S. K. Patra,
 - [14] S. K. Patra, R. K. Gupta and W. Greiner, *Mod. Phys. Lett. A* **12**, 1727 (1997); Tapas Sil, S.K. Patra, B.K. Sharma, M. Centelles, and X. Viñas, *Phys. Rev. C* **69**, 044315 (2004).
 - [15] Yu. Ts. Oganessian, A. S. Iljinov, A. G. Demin, S. P. Tretyakova, *Nucl. Phys. A* **239**, 353 (1975).
 - [16] Yu. Ts. Oganessian et al, *Phys. Rev. C* **74**, 044602 (2006).
 - [17] Yu. Ts. Oganessian et al, *Phys. Rev. C* **83**, 054315 (2011).

- [18] J. V. Kratz, *Heavy Elements and Related New Phenomena*, edited W. Greiner and R. K. Gupta (World Scientific, Singapore, 1999), Vol. I, Chap. 4, p. 129.
- [19] A. Sobiczewski and K. Pomorski, Prog. in Part. and Nucl. Phys. **58**, 292 (2007).
- [20] J. Decharge, J. -F. Berger, K. Dietrich and M. S. Weiss, Phys. Lett. **B 451**, 275 (1999); J. Decharge, J. -F. Berger, M. Girod and K. Dietrich, Nucl. Phys. **A 716**, 55 (2003).
- [21] P.-G. Reinhard and H. Flocard, Nucl. Phys. A **584**, 467 (1995).
- [22] G. A. Lalazissis, S. Karatzikos, R. Fossion, D. Pena Arteaga, A. V. Afanasjev, and P. Ring, Physics Letters B, **671**, 36 (2009).
- [23] J.R. Stone, J.C. Miller, R. Koncewicz, P.D. Stevenson, and M.R. Strayer, Phys. Rev. C **68** 034324 (2003).
- [24] M. Bender, P. -H. Heenen and P. -G. Reinhard, Rev. Mod. Phys. **75**, 121 (2003).
- [25] M. Dutra, O. Lourenco, J. S. Sa Martins, A. Delfino, J.R. Stone and P. D. Stevenson, Phys. Rev. C **85** 035201 (2012).
- [26] J. R. Stone and P. G. Reinhard, Prog. Part. Nucl. Phys. **58**, 587 (2007).
- [27] E. Chabanat, P. Bonche, P. Hansel, J. Mayer, R. Schaeffer, Nucl. Phys. **A 627** 710 (1997).
- [28] E. Chabanat, P. Bonche, P. Haensel, J. Meyer, and R. Schaeffer, Nucl. Phys. **A635** 231 (1998).
- [29] B.A. Brown, Phys. Rev. **C58** 220 (1998).
- [30] B. D. Serot, Rep. Prog. Phys. **55**, 1855 (1992).
- [31] Y. K. Gambhir, P. Ring, and A. Thimet, Ann. Phys. (N.Y.) **198**, 132 (1990).
- [32] P. Ring, Prog. Part. Nucl. Phys. **37**, 193 (1996).
- [33] B. D. Serot and J. D. Walecka, Adv. Nucl. Phys. **16**, 1 (1986).
- [34] P. K. Panda, S. K. Patra, J. Reinhardt, J. A. Maruhn, H. Stocker, and W. Greiner, Int. J. Mod. Phys. E **6**, 307 (1997).
- [35] G. Audi, A.H. Wapstra and C. Thibault, Nucl. Phys. A, **729**, 337 (2003).
- [36] P. Moller, J. R. Nix, W. D. Myers, and W. J. Swiatecki, At. Data and Nucl. Data Tables **59**, 185 (1995).
- [37] P. Moller, J. R. Nix, and K.-L. Kratz, At. Data and Nucl. Data Tables **66**, 131 (1997).
- [38] J. P. Maharana, Y. K. Gambhir, J. A. Sheikh and P. Ring, Phys. Rev. C **46**, R1163 (1992).
- [39] S. K. Patra, and C. R. Praharaj, Phys. Rev. C **47**, 2978 (1993).
- [40] F. Sarazin et. al., Phys. Rev. Lett. **84**, 5062 (2000).
- [41] J. L. Egido, L. M. Robledo, R. R. Rodriguez-Guzman, Phys. Rev. Lett. **93**, 282502 (2004).
- [42] J.A.Wheeler (prior to 1950) unpublished.

- [43] H. A. Wilson, Phys. Rev. **69**, 538 (1946).
- [44] P. J. Siemens and H. A. Bethe, Phys. Rev Lett. **18**, 704(1967).
- [45] C. Y. Wong, Phys. Lett. B **41**, 451 (1972).
- [46] K. T. R. Davies, C. Y. Wong and S. J. Krieger, Phys. Lett. B **41**, 455 (1972).
- [47] J. Decharge et. al., Nucl. Phys. A **716**, (2003).
- [48] M. Grasso et. al., Phys. Rev. C **79**, 034318 (2009).
- [49] W. Y. Zao. et. al., Chin. Phys. Lett, **28**, 102101 (2011).
- [50] E. Khan. et. al., Nucl. Phys. A, **800**, 37 (2008).
- [51] S. K. Patra and C. R. Praharj, J. Phys. G **23**, 939 (1997).
- [52] V.E. Viola, Jr. and G.T. Seaborg, J. Inorg. Nucl. Chem. **28**, 741(1966).
- [53] P. Moeller, J.R. Nix, and K.L. Kratz, At. Data Nucl. Data Tables **66**, 131(1997).
- [54] W.D. Myers and W.J. Swiatecki, Phys. Rev. C **58**, 3368(1998).
- [55] S. K. Singh and S. K. Patra (In progress).